

## TURF-TYPE TALL FESCUE LAWN CALENDAR

	<b>B B</b> est month			Р		Possible Month		Μ		Marginal month		
I 7 8 5	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
MOWING HEIGHT: 2"-3" ; raise 1/2" higher during dry periods. Remove no more than 1/3 total height at one time.	OK	ОК	ОК	OK	OK	OK	OK	ОК	ОК	ОК	OK	OK
WATER: 1 inch per week if no rainfall.	ОК	ОК	ОК	ОК	ОК	ОК	ОК	ОК	ОК	ОК	ОК	ОК
<b>FERTILIZATION:</b> Use any turf fertilizer. Follow label rate.		В	Р	В					В	Р	В	
<b>IDEAL pH RANGE: 5.5-6.5</b> Use dolomitic lime per soil test recommendation. Can be applied at any time.												
<b>AERATION:</b> Use a core aerator during active growth season.	м	М	Р	В	м				Р	В	Р	м
<b>DETHATCH:</b> Generally not necessary.												
<b>SEEDING - NEW LAWN:</b> 5 - 8 lb. of seed per 1000 sq. feet.		Р	Р	М	М				Р	В	Р	
<b>OVERSEEDING - ESTABLISHED LAWN:</b> 3 - 5 lb. seed per 1000 sq. feet		В	Р	Р	М				Р	В	Р	
<b>SODDING</b> Fescue sod is not available in all months of the year.	В	В	В	Р	М				Р	В	В	В
WEED CONTROL: Read product label carefully to determine which weeds are controlled and on which grasses the product can be used.												
<b>Spring pre-emergent*</b> (prevents crabgrass, goosegrass and other summer weed seeds from sprouting)		В	В	м								
Broadleaf post-emergent* (spot spray to kill broadleaf plants like chickweed, wild violet, dandelion, wild onion, etc)	ОК	ОК	ОК	ОК	ОК	ОК	ОК	ОК	ОК	ОК	ОК	ОК
<b>Grassy weed post-emergent</b> * (kills grassy weeds like crabgrass, dallisgrass, etc. Do not apply to drought-stressed fescue.)					Р	Р	М					
Winter pre-emergent* (prevents chickweed and other winter weed seeds from sprouting)									В	Р		

\* Read weed control product labels carefully. Some products can not be applied to lawns that will be seeded within a few weeks. Do not apply herbicides to newly sprouted fescue until it has been mowed four times.